

GET THE FACTS

Available Outpatient COVID-19 Treatments

Sick with COVID-19 symptoms?

Treatments are now available for nonhospitalized adults and some children with COVID-19.

These prescription treatments:

- ✓ are FDA approved or authorized.
- ✓ decrease serious disease and hospitalizations.
- ✓ work best when used early.

For use within 5 days of first symptoms:

NIRMATRELVIR/RITONAVIR
(PAXLOVID™)



2x
for



2-3 pills twice a day for 5 days

For use within 7 days of first symptoms:

REMDESIVIR
(VEKLURY®)

1 IV infusion (i.e., delivered through the vein) per day for 3 days



1x
for



MOLNUPIRAVIR
(LAGEVRIO®)

4 pills twice a day for 5 days



2x
for



BEBTELOVIMAB
(MONOCLONAL ANTIBODY)

Single IV infusion over 30 seconds (followed by 1 hour of monitoring)



1x
for



For use within 8 days of first symptoms:

**HIGH-TITER
CONVALESCENT PLASMA**

Single IV transfusion over 60 minutes



1x
for



Good to Know

- ⚠ There is only a short time window when these treatments can be used. So at first sign of illness, get tested for COVID-19 and contact your health care provider or visit covid.gov.
- ⚠ There may be issues with drug interactions using nirmatrelvir/ritonavir (Paxlovid™) with other common medicines, so check with your health care providers.
- ⚠ Molnupiravir should NOT be given to children or pregnant people, and:
 - Women should use contraception during the course.
 - Men should use contraception during the course and for 3 months after.
- ⚠ These medications are not a substitute for vaccination or other methods to prevent COVID-19.



For more information on therapeutics, visit:
COVID19LearningNetwork.org

COVID-19 Real-Time Learning Network

Brought to you by **CDC** and **NIH**

